

An Overview of the Functional Linear Model

We want to see where these lectures
on the functional linear model will go.

A functional analysis of . . .

A scalar response and . . .

A functional response . . .

What exactly makes a . . .

Home Page

Title Page

◀ ▶

◀ ▶

Page 1 of 13

Go Back

Full Screen

Close

Quit

- Four functional linear models for the daily weather data.
- A functional ANOVA for precipitation.
- Predicting total annual precipitation from the temperature profile.
- Predicting today's precipitation from today's temperature.
- Predicting the entire year's precipitation from the year's temperature profile.
- A short-term feed-forward model for precipitation.
- A more general perspective.
- Predicting precipitation dynamics: a differential equation
- The idea of a linear model reviewed.

A functional analysis of . . .

A scalar response and . . .

A functional response . . .

What exactly makes a . . .

Home Page

Title Page

◀ ▶

◀ ▶

Page 2 of 13

Go Back

Full Screen

Close

Quit

The average Canadian weather data

- 35 Canadian weather stations selected to cover the country.
- Daily temperatures (0.1 degrees Celsius) and precipitations (0.1 mm) averaged over the years 1960 to 1994. (Feb 29th combined with Feb. 28th).
- Canada divided into Atlantic, Continental, Pacific and Arctic weather zones.

Home Page

Title Page

◀ ▶

◀ ▶

Page 3 of 13

Go Back

Full Screen

Close

Quit



A functional analysis of ...

A scalar response and ...

A functional response ...

What exactly makes a ...

[Home Page](#)

[Title Page](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

Page 4 of 13

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)

1. A functional analysis of variance

- Does the precipitation profile vary from one weather zone to another?
- We have a number N_g of weather stations in each climate zone $g = 1, \dots, 4$, and
- the model for the m th temperature function in the g th group, indicated by Prec_{mg} , is

$$\text{Prec}_{mg}(t) = \mu(t) + \alpha_g(t) + \epsilon_{mg}(t).$$

- $\mu(t)$ is the grand mean function, summarizing precipitation for all of Canada.
- $\alpha_g(t)$ is the functional effect of being in weather zone g
- In order to fix zone effects, we require that

$$\sum_g \alpha_g(t) = 0 \text{ for all } g$$

A functional analysis of ...

A scalar response and ...

A functional response ...

What exactly makes a ...

Home Page

Title Page

◀◀ ▶▶

◀ ▶

Page 5 of 13

Go Back

Full Screen

Close

Quit

2. A scalar response and a functional independent variable

- The response is the log total annual precipitation

$$\text{PrecTot}_i = \int_0^{365} \text{Prec}_i(t) dt$$

- The model is

$$\log(\text{PrecTot}_i) = \alpha + \int_0^{365} \text{Temp}_i(s) \beta(s) ds + \epsilon_i.$$

- But here we have a real problem. How to avoid over-fitting the 35 scalar observations?
- We'll use regularization or roughness penalties on the estimated regression functions.

A functional analysis of ...

A scalar response and ...

A functional response ...

What exactly makes a ...

Home Page

Title Page

◀ ▶

◀ ▶

Page 6 of 13

Go Back

Full Screen

Close

Quit

3. A functional response and a functional independent variable

This is a big topic, and breaks down into several useful special versions.

[Home Page](#)[Title Page](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 7 of 13](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

3.1. The concurrent functional model

- We might only use the temperature at the same time $s = t$ because we imagine that precipitation now depends only on the temperature now.
- Our model is

$$\text{Prec}_i(t) = \alpha(t) + \text{Temp}_i(t)\beta(t) + \epsilon_i(t)$$

- We might call this model *concurrent* or *point-wise*.
- Should we use regularization to force β to be smooth in t ?

[Home Page](#)[Title Page](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)

Page 8 of 13

[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

3.2. The annual or total model

- We may prefer to allow for temperature influence on $\text{Prec}(t)$ to extend over the whole year.
- The model expands to become

$$\text{Prec}_i(t) = \alpha(t) + \int_0^{365} \text{Temp}_i(s) \beta(s, t) ds + \epsilon_i(t)$$

- The value $\beta(s, t)$ determines the impact of temperature at time s on precipitation at time t .
- We need roughness penalties for variation in both s and t

[Home Page](#)[Title Page](#)[◀](#) [▶](#)[◀](#) [▶](#)

Page 9 of 13

[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

3.3. The limited-term feed-forward model

- it may be that what counts is whether the temperature has been falling rapidly up to time t . The model expands to

$$\text{Prec}_i(t) = \alpha(t) + \int_{t-\delta}^t \text{Temp}_i(s) \beta(s, t) ds + \epsilon_i(t)$$

- Here δ is the time lag over which we use temperature information.
- Now β is only defined over the somewhat complicated trapezoidal domain: $t \in [0, 365], t - \delta \leq s \leq t$.

[Home Page](#)
[Title Page](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Page 10 of 13](#)
[Go Back](#)
[Full Screen](#)
[Close](#)
[Quit](#)

3.4. The local influence model

- Finally, we may open up the model to allow integration over s within a t -dependent set Ω_t .
- The model may therefore be

$$\text{Prec}_i(t) = \alpha(t) + \int_{\Omega_t} \text{Temp}_i(s) \beta(s, t) ds + \epsilon_i(t)$$

[Home Page](#)[Title Page](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 11 of 13](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

3.5. What about predicting derivatives?

- When the response is a derivative, then there is the potential for the function itself to be a useful covariate.
- The concurrent linear model

$$DPrec_i(t) = Prec_i(t)\beta(t) + \epsilon_i(t)$$

is a *homogeneous first order linear differential equation* in precipitation.

- If we also include an influence of temperature,

$$DPrec_i(t) = Prec_i(t)\beta_0(t) + Temp_i(t)\beta_1(t) + \epsilon_i(t),$$

the equation is said to be *forced* or *nonhomogeneous*.

[Home Page](#)
[Title Page](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)

Page 12 of 13

[Go Back](#)
[Full Screen](#)
[Close](#)
[Quit](#)

4. What exactly makes a model linear?

- We see that the functional linear model has a lot more variants than it's multivariate cousin.
- We will need to look at a definition of a linear model that encompasses these models and many others.

[Home Page](#)[Title Page](#)[◀](#) [▶](#)[◀](#) [▶](#)[Page 13 of 13](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)